

Notes on the Lamiaceae of China

Li Xi-wen (Li Hsi-wen)

Kunming Institute of Botany, Academia Sinica, Kunming, Yunnan 650204, People's Republic of China

ABSTRACT. Nomenclatural changes, affecting *Marmoritis* and *Scutellaria*, are made for the upcoming *Flora of China*.

Hedge (1990) resuscitated the generic name *Marmoritis* "after almost 150 years in synonymy." It replaces the previously accepted *Phyllophyton*. In this same treatment Hedge made the new combination *Marmoritis nivalis* (Jacquemont ex Benth.) Hedge and recognized *Marmoritis rotundifolia* Bentham for the flora area he was considering. In order to reflect the current taxonomic status for species formerly placed in *Phyllophyton*, the following nomenclatural changes are made. The taxonomic treatment of the following taxa, along with *Marmoritis nivalis* and *M. rotundifolia*, will appear in the *Flora of China* (Li & Hedge, in press).

***Marmoritis complanatum* (Dunn) H. W. Li, comb. nov.** Basionym: *Nepeta complanata* Dunn, Notes Roy. Bot. Gard. Edinburgh 8: 122. 1913. *Glechoma complanata* (Dunn) Turrill, Bot. Soc. Exch. Club Brit. Isles 1919, 5: 695. 1920. *Phyllophyton complanatum* (Dunn) Kudô, Mem. Fac. Sci. Taihoku Imp. Univ. 2: 225. 1929. *Pseudolophanthus complanatus* (Dunn) Levin, Trudy Bot. Inst. Akad. Nauk S.S.R., Ser. 1, Fl. Sist. Vyssh. Rast. 5: 296. 1941. TYPE: China. NW Yunnan: scree plant, 16,000 ft., July 1911, F. Kingdon Ward 138 (holotype, E not seen).

Dracocephalum rockii Diels, Notizbl. Bot. Gart. Berlin-Dahlem 9: 1030. 1926. TYPE: China. Yunnan Province: Mountains of Moting, NW of the Yangtze-Mekong watershed, June 1923, J. F. Rock 10327 (holotype, presumably B, not seen; isotype, US).

***Marmoritis decolorans* (Hemsley) H. W. Li, comb. nov.** Basionym: *Nepeta decolorans* Hemsley, Hooker's Icon. Pl. 25: t. 2470. 1896. *Glechoma decolorans* (Hemsley) Turrill, Bot. Soc. Exch. Club Brit. Isles 1919, 5: 695. 1920. *Phyllophyton decolorans* (Hemsley) Kudô, Mem. Fac. Sci. Taihoku Imp. Univ. 2: 225. 1929, "decorans." *Pseudolophanthus decolorans* (Hemsley) Levin, Trudy Bot. Inst. Akad. Nauk S.S.R., Ser. 1, Fl. Sist. Vyssh. Rast. 5: 296. 1941. TYPE: Hab. Central Tibet: Gooring Valley, 30°12'N, 90°25'E, at about 16,500 ft., St. George Littledale, July and Aug. 1895 (holotype, K not seen).

***Marmoritis pharicus* (Prain) H. W. Li, comb. nov.** Basionym: *Nepeta pharica* Prain, J. Asiat. Soc. Bengal, Pt. 2, Nat. Hist. 59: 306. 1891. *Glechoma pharica* (Prain) Turrill, Bot. Soc. Exch. Club Brit. Isles 1919, 5: 695. 1920. *Phyllophyton pharicum* (Prain) Kudô, Mem. Fac. Sci. Taihoku Imp. Univ. 2: 225. 1929. *Pseudolophanthus pharicus* (Prain) Kuprianova, Bot. Zhurn. S.S.R. 33: 235. 1948. TYPE: Eastern Tibet, Phari, between Phari and Lhasa, Lama Uyen Gyatsko 106 (holotype, not seen).

***Scutellaria yangbiensis* H. W. Li, nom. nov.** Replaced name: *Scutellaria urticifolia* C. Y. Wu & H. W. Li, Fl. Yunnan. 1: 545. 1977, non Juzepczuk & Vvedensky, Fl. URSS 20: 510. 1954. TYPE: Zhongdian Xian Expedition, NW Yunnan, 63-4020 (holotype, KUN).

Acknowledgments. The assistance of Bryan Dutton and Benito Tan is gratefully acknowledged.

Literature Cited

- Hedge, I. 1990. Labiateae. *Flora of Pakistan* 192: 119.
Li, X. W. & I. Hedge. Lamiaceae. *Flora of China*, vol. 17, Science Press & Missouri Bot. Gard. (In press.)